arXiv:1606.07023v1 [math.HO] 21 Jun 2016

A SPECIAL THEOREM RELATED TO THE FAGNANO'S PROBLEM

JUN LI

ABSTRACT. A special theorem related to the Fagnano's problem is proved and an example of the theorem is shown in a golden rectangle.

1. A SPECIAL THEOREM RELATED TO THE FAGNANO'S PROBLEM

The Fagnano's problem (see, e.g., [1]) is an optimization problem that was first stated by Giovanni Fagnano in 1775:

Problem 1.1. *For a given acute triangle $\triangle ABC$, determine the inscribed triangle of minimal perimeter.*

The answer to Problem 1.1 is that the orthic triangle of $\triangle ABC$ has the smallest perimeter. Here, we have a special Theorem 1.2 related to the problem.



FIGURE 1. A special theorem related to the Fagnano's problem

Theorem 1.2. *In Figure 1, Suppose $\triangle ABC$ is an acute-angled triangle, then, the of smallest perimeter triangle $\triangle DEF$ can be inscribed in $\triangle ABC$ is a right triangle if and only if $\triangle ABC$ has only one $\frac{\pi}{4}$ angle. In addition, the right angle $\angle E$ and the $\frac{\pi}{4}$ angle are opposite angles, which means here, $\angle B = \frac{\pi}{4}$.*

Proof. According to the answer above, the $\triangle DEF$ is the orthic triangle of $\triangle ABC$, and it's also known that the incenter of the orthic triangle $\triangle DEF$ is the orthocenter of $\triangle ABC$ (see, e.g., [2]). Now, suppose $\triangle DEF$ is a right triangle with $\angle E = \frac{\pi}{2}$, then in $\triangle DEF$, we have

$$\angle D + \angle E + \angle F = \pi$$

$$\angle DFC = \angle CFE, \angle FDA = \angle ADE, \angle FEB = \angle BED = \frac{\pi}{4}$$

Date: 2016.6.

then we get

$$\angle CFE + \angle ADE = \frac{\pi}{4}$$

and in the quadrilateral $BDEF$, we have

$$\angle B + \angle E + \angle BFE + \angle BDE = 2\pi$$

$$\angle BFE = \angle BFC + \angle CFE = \frac{\pi}{2} + \angle CFE$$

$$\angle BDE = \angle BDA + \angle ADE = \frac{\pi}{2} + \angle ADE$$

with $\angle E = \frac{\pi}{2}$, we conclude $\angle B = \frac{\pi}{4}$, and obviously, an acute-angled triangle can only have one $\frac{\pi}{4}$ angle, and also, it's easy to see that the right angle $\angle E$ and $\angle B$ are opposite angles. □



FIGURE 2. An example of the special theorem in a golden rectangle

Next, we show an example of the special Theorem 1.2 in a golden rectangle[5][4, p. 274][6, p. 115].

Example 1.3. *In Figure 2, $ABCD$ is a golden rectangle with $AB = 1$, $BC = \phi$, made up of a unit square $ABEF$ and a small golden rectangle $FECD$, then the of smallest perimeter triangle $\triangle GHE$ that can be inscribed in the acute-angled triangle $\triangle BFC$ is a right triangle having sides proportional to* $(1, 2, \sqrt{5})$[3].

Proof. It's easy to see that $\triangle BFC$ is an acute-angled triangle with $\angle B = \frac{\pi}{4}$, and according to Theorem 1.2, the orthic triangle $\triangle GHE$ is a right triangle with $\angle H = \frac{\pi}{2}$. Since $\triangle BGC \sim \triangle BEF$, we have $\frac{BG}{BC} = \frac{BE}{BF} = \frac{1}{\sqrt{2}}$ and get $BG = \frac{\phi}{\sqrt{2}}$, then apply the law of cosines to $\triangle GBE$, we have

$$GE^2 = BG^2 + BE^2 - 2BG \cdot BE \cos \angle B$$

and get $GE = \sqrt{\frac{1+\phi^2}{2\phi^2}}$. Using the similar method, we can get $HE = \sqrt{\frac{2}{1+\phi^2}}$, hence $\frac{GE}{HE} = \frac{\sqrt{5}}{2}$. □

REFERENCES

[1] Fagnano's problem, *Wikipedia, The Free Encyclopedia*, https://en.wikipedia.org/wiki/Fagnano_problem.
[2] Altitude (triangle), *Wikipedia, The Free Encyclopedia*, https://en.wikipedia.org/wiki/Altitude_(triangle)#Orthic_
[3] C. Radin, The pinwheel tilings of the plane, *Annals of Mathematics.* **139.3** (1994) 661–702.
[4] T. Koshy, *Fibonacci and Lucas Numbers with Applications*, A Wiley-Interscience Publication, 2001.
[5] M. Bicknell and V. E. Hoggatt, Golden triangles, rectangles, and cuboids, *The Fibonacci Quarterly*, **7.1** (1969) 73–91.
[6] A. S. Posamentier and I. Lehmann, *The Fabulous Fibonacci Numbers*, Prometheus Books, 2007.

Mathematics Subject Classification (2010). 51M04, 11B39
Keywords. Fagnano's problem, Golden rectangle

SCHOOL OF SCIENCE, JIANGXI UNIVERSITY OF SCIENCE AND TECHNOLOGY, GANZHOU, 341000, CHINA.
E-mail address: junli323@163.com